

REVIEW ARTICLE

***Vernonia amygdalina* Del: A Mini Review**

Suleiman Danladi^{1*}, Mohammed Alkassim Hassan¹, Idris Aliyu Masa'ud², Umar Idris Ibrahim³

¹Department of Pharmaceutical and Medicinal Chemistry, Faculty of Pharmaceutical Sciences, Bayero University, Kano, Nigeria.

²Department of Pharmacognosy and Herbal Medicine, Faculty of Pharmaceutical Sciences, Bayero University, Kano, Nigeria.

³Department of Clinical Pharmacy and Pharmacy Practice, Faculty of Pharmaceutical Sciences, Bayero University, Kano, Nigeria.

*Corresponding Author E-mail: danladisuleimen@gmail.com

ABSTRACT:

Phytomedicines are medicinal products of plant origin that have been used since ancient times till date for the prophylaxis and treatment of various ailments. *Vernonia amygdalina* Del (Asteraceae) is a perennial shrub that is widely distributed in tropical parts of Africa. It is popularly known as bitter leaf and consumed as a vegetable and for medicinal purposes. The plant has been used in folkloric medicine for the treatment of several diseases owing to its diverse pharmacological applications. The plant has low toxicity profile; thus, making it a potential drug candidate. Scientific investigations have shown that, *V. amygdalina* Del possesses potent pharmacological activities such as antidiabetic, anthelmintic, antiplasmodial, antimicrobial, antioxidant and antianaemic activities. Moreover, *V. amygdalina* Del contains various phytochemical compounds which may be associated with its potent pharmacological properties. The present study is a review of the therapeutic benefits as well as phytochemicals isolated from *V. amygdalina* Del.

KEYWORDS: *Vernonia amygdalina*, phytomedicine, pharmacological-activity, phytochemical.

INTRODUCTION:

Herbal medicines, otherwise called phytomedicines, are medicinal products of plants or plant parts such as roots, leaves, flowers, barks, seeds and fruits which are used to treat diseases and to improve health. They are used as complementary or alternative medicines; in the prophylactic treatment and alleviation of diseases and their symptoms.¹ Medicinal plants play a vital role in the treatment of various diseases as evident by their long history of use. They have been used in both ancient and recent times by all culture for curing several diseases²

Vernonia amygdalina Del is among the medicinal plants used for the treatment of various diseases in different parts of the world. It has been reported that *V. amygdalina* Del plant is used for the treatment of diabetes, yellow fever, dysentery, constipation, malaria and stomach ache in Africa and Asia.^{2,3} Moreover, the plant is widely used in Africa as a source of food as vegetable and culinary herb in soup.⁴

V. amygdalina Del belongs to the family Asteraceae. It is widely distributed throughout Tropical Africa and cultivated as food supplement in West Africa including Nigeria.^{2,5} The common name of *V. amygdalina* Del is 'bitter leaf', locally known as 'Shuwaka' in Hausa Language and 'Ewuro' in Yoruba Language.⁶

In this research various activities as well as phytochemical constituents of *V. amygdalina* Del were reviewed. The data were searched from online sources; Pubmed, googlescholar and science direct. The key word used for searching information was *V. amygdalina* Del.

Taxonomy:

Vernonia amygdalina Del. is a small tree growing up to 3 m high. The picture of *V. amygdalina* Del in its natural habitat is shown in figure 1.0. It grows throughout the African Tropics⁷.



Figure 1.0: *Vernonia amygdalina* Del plant

V. amygdalina commonly called bitter leaf is a perennial shrub belonging to the family *Asteraceae*. It grows throughout Tropical Africa.⁸Its taxonomic classification⁶ is presented in Table 1.0.

Table 1.0: Taxonomic classification of *V. amygdalina*Del

Kingdom	Plantae
Class	Angiospermae
Order	Asterales
Family	Asteraceae
Genus	<i>Vernonia</i>
Species	<i>amygdalina</i> Del

Phytochemistry of *V. amygdalina* Del:

Qualitative phytochemical screening of *V. amygdalina* Del have detected heavy presence of polyphenols and moderate presence of alkaloids, saponins, flavonoids and steroids⁵. Phytochemical study of the leaves of *V. amygdalina* Del showed that, the plant contains two sesquiterpene lactones; vernolide and vernodalol.⁷ Vernoniosides D and Vernoniosides E were isolated from the leaves of *V. amygdalina* Del⁹.







Figure 1: Some compounds isolated from *V. amygdalina*

Antidiabetic activity:

V. amygdalina Del, demonstrated positive antidiabetic activity when administered through various routes of administration. Intraperitoneal administration of aqueous leaf extract of *V. amygdalina*, Del, produced hypoglycaemic effects which was comparable to tolbutamide. Additionally, when aqueous leaf extract of *V. amygdalina*, was given to normoglycaemic rabbits, the fasting blood sugar reduced from 96 mg% to 48 mg% in 4 h.¹⁰ Similarly, administration of the leaf extract via gastric intubation to alloxan induced rats provided significant reduction of blood and serum glucose.¹¹ *V. amygdalina* Del has been reported to contain high amount of polyphenols.¹¹ Accordingly, phenolic extract of *V. amygdalina* Del was found to have significant effect on the key enzymes linked to type-2 diabetes thereby inhibiting the activities of α -amylase and α -glucosidase *in vitro* in a concentration-dependent manner.¹² Okolie *et al.* (2008)¹³ assessed the antidiabetic effect of *V. amygdalina* Del in human subjects using standard laboratory testing (fasting blood glucose, glucose tolerance test and postprandial blood glucose). It was found that *V. amygdalina* Del caused significant reductions in blood glucose levels almost postprandial time points and for area-under-curve.¹³

The polyherbal combination of *V. amygdalina* Del and *Azadirachta indica* constituents gave a synergistic antidiabetic effect in diabetic rat.³ In another study,

combination of *V. amygdalina* Del and *Gongronema latifolium* provided significant reduction of serum glucose in streptozotocin (STZ)-induced diabetic Wistar rats. This effect was comparable to that of insulin.¹⁴

Atangwho *et al.* (2010)¹¹ suggested that, the mechanism of antidiabetic activity of *V. amygdalina* Del may be due to the regeneration of pancreatic β -cell.

Anthelmintic activity:

The acetone extract of *V. amygdalina* Del was found to inhibit parasitic larva of *Haemonchus contortus*. It was also reported to inhibit nematode eggs and nematode larvae¹⁵. In another study, fifty West African dwarf goats were used to compare the anthelmintic activity of some standard drugs namely albendazole, levamisole and ivermectin with that of the aqueous extract of bitter leaf using the Faecal Egg Count Reduction (FECR) assay. The aqueous extract demonstrated highest FECR (100%) compared to the standard drugs (albendazole, 99%; levamisole, 96%; ivermectin, 96%) against the nematodes (*Haemonchus contortus*, *Trichostrongylus* spp., *Ostertagia ostertagi*, *Oesophagostomum* spp., *Chabertia* sp., *Strongyloides*, and *Paramphistomum* spp.) tested.¹⁶

Antiplasmodial activity:

The aqueous and ethanolic leaf extracts of *V. amygdalina* Del were analysed *in vivo* (in rats) and *in vitro* (cell lines) using bioassay guided fractionation for the isolation of molecules with antimalarial activity. Both extracts demonstrated antiplasmodial activities at both sexual and asexual blood stages of the parasite.¹⁷ *V. amygdalina* Del leaf was reported to have antiplasmodial activity *in vitro* which was attributed to its phytochemical constituents¹⁸.

Antimicrobial activity:

Vernolide and vernodalol; the chemical compounds isolated from the leaves of *V. amygdalina* Del demonstrated a significant bactericidal activity against Gram positive bacteria. However, they failed to show efficacy against the Gram negative strains.⁷ Moreover, vernolide and vernodalol showed high and moderate activity respectively against some fungal species. In contrast, they were ineffective against *Fusarium oxysporum*⁷.

Antioxidant activity:

Methanolic extract of *V. amygdalina* Del prevented lipid peroxidation induced by tert-butyl hydroperoxide.¹⁹ It was also reported that, the methanolic extract of *V. amygdalina* exhibited significant free radical scavenging and antioxidant activities²⁰.

Antianaemic activity:

Methanolic extract of *V. amygdalina* Del lowered erythrocytes lysis induced by tert-butyl hydroperoxide.¹⁹ Therefore, anti-haemolytic effect of *V. amygdalina* Del suggests that the plant can be used as antianaemic agent.

Effect on fertility:

The reproductive effect of saponin extract of *V. amygdalina* Del studied on male Wistar rats at different doses for 14 days showed that sperm cell motility and concentration were enhanced (at higher doses) without a reduction in the percentage of morphologically abnormal sperm cells²¹.

Immunomodulatory effect:

V. amygdalina Del has been studied for its modulatory effect on the immune system. In one study, the immunomodulatory effect of the leaf extract of the plant, Immunace® and a combination of both were observed in human immunodeficiency virus (HIV) patients on Anti-Retroviral Therapy (ART). A combination of the extract with Immunace® improved the CD₄⁺ counts and packed cell volume (PCV) while reducing the development of skin rashes of patients when compared to control (patients on ART only).²² Similarly, Ezeonu *et al.* (2016)²³ analysed the leaf extract of *V. amygdalina* Del for its immunological effect in rats and reported a remarkable improvement on CD₄⁺ counts, haematological parameters and lipid profiles.

Cardiovascular Activity:

The cardiovascular activity of the aqueous leaf extract of *V. amygdalina* Del was investigated in rabbits fed with atherogenic diet. There was no significant improvement of lipid profile when the bitter leaf-atherogenic diet fed rabbits were compared to control.²⁴

Safety and Toxicities Profile:

Olujimi *et al.* (2017)²⁵ analyzed concoctions containing ten medicinal plants (including bitter leaf) with potential antidiabetic activity for the presence of toxic elements. They were found to be safe from the risk of the toxic metals and trace elements investigated. There was no death or any physical pathology observed at doses of up to 600 mg/kg when aqueous extract of *V. amygdalina* Del was administered to rats.²⁶ The histopathology and microscopy of the liver, kidney, small intestine, lung and heart did not show any significant damage except for mild congestion in the kidney after oral administration of *V. amygdalina* Del extract in rats. Biochemical analysis revealed that *V. amygdalina* Del has the potential to elevate some liver enzymes (ALT, AST and ALP) and cholesterol in bitter leaf extract-treated rats.²⁶ Interestingly, in an *in vitro* study against plasmodium parasite *V. amygdalina* Del leaf extract has been shown to have no sign of toxicity¹⁸.

CONCLUSION:

V. amygdalina Del is a versatile plant that possesses several prophylactic and therapeutic potentials. This review highlighted many of the pharmacological activities of the plant. Interestingly *V. amygdalina* Del leaf was reported to have low toxicity profile, thus making it a potential drug candidate. However, further researches are required to standardize extracts of the plant and to unravel further medicinal and nutraceutical properties of the plant particularly on the root, stem and bark.

CONFLICT OF INTEREST:

No conflict of interest.

REFERENCES:

- Calixto JB, Santos AR, Filho VC, Yunes RA. A review of the plants of the genus *Phyllanthus*: their chemistry, pharmacology, and therapeutic potential. *Medicinal research reviews*. 1998 Jul 1;18 (4):225-58.
- Adebite AE, Sanyaolu EB. Cytotoxicity testing of aqueous extract of bitter leaf (*Vernonia amygdalina* Del.) using the *Allium cepa* chromosome aberration assay. *Scientific Research and Essays*. 2009 Nov 30;4 (11):1311-4.
- Ebong PE, Atangwho IJ, Eyong EU, Egbung GE. The antidiabetic efficacy of combined extracts from two continental plants: *Azadirachta indica* (A. Juss)(Neem) and *Vernonia amygdalina* (Del.)(African bitter leaf). *American Journal of Biochemistry and Biotechnology*. 2008;4 (3):239-44.
- Nwosu SI, Stanley HO, Okerentugba PO. Occurrence, types and location of calcium oxalate crystals in *Vernonia amygdalina* Del (Asteraceae) *Int. J. Sci. Nat.*. 2013; 4(3):533-537
- Atangwho IJ, Ebong PE, Eyong EU, Williams IO, Eten MU, Egbung GE. Comparative chemical composition of leaves of some antidiabetic medicinal plants: *Azadirachta indica*, *Vernonia amygdalina* and *Gongronema latifolium*. *African Journal of Biotechnology*. 2009;8(18).
- Farombi EO, Owoeye O. Antioxidative and chemopreventive properties of *Vernonia amygdalina* and *Garcinia biflavonoid*. *International Journal of Environmental Research and Public Health*. 2011;8(6):2533-55.
- Erasto P, Grierson DS, Afolayan AJ. Bioactive sesquiterpene lactones from the leaves of *Vernonia amygdalina*. *Journal of Ethnopharmacology*. 2006 ;106 (1):117-20.
- Ijeh II, Ejike CE. Current perspectives on the medicinal potentials of *Vernonia amygdalina* Del. *Journal of Medicinal Plants Research*. 2011 Apr 4;5 (7):1051-61.
- Igile G, Olenszek W, Jurzysta M, Aquino R, de Tommasi N, Pizzi C. *Vernonia amygdalina* D and E, two novel saponins from *Vernonia amygdalina*. *Journal of Natural Products*. 1995 Sep;58 (9):1438-43.
- AkahPA, Okafor CL. Blood sugar lowering effect of *Vernonia amygdalina* Del, in an experimental rabbit model. *Phytotherapy Research*. 1992, 6 (3):171-173.
- Atangwho IJ, Ebong PE, Egbung GE, Obi AU. Extract of *Vernonia amygdalina* Del.(African bitter leaf) can reverse pancreatic cellular lesion after alloxan damage in the rat. *Australian Journal of Basic and Applied Sciences*. 2010;4(5):711-6.
- SaliuJA, AdemiluyiAO, AkinyemiAJ, Oboh G. *In Vitro* antidiabetes and antihypertension properties of phenolic extracts from bitter leaf (*Vernonia amygdalina* DEL.) *Journal of Food Biochemistry*. 2012; 36 (5):569-576.
- Okolie UV, Okeke CE, Oli JM, Ehiemere IO. Hypoglycemic indices of *Vernonia amygdalina* on postprandial blood glucose concentration of healthy humans. *African Journal of Biotechnology*. 2008;7 (24).
- Akpaso MI, Atangwho IJ, Akpantah A, Fischer VA, Igiri AO, Ebong PE. Effect of Combined Leaf Extracts of *Vernonia amygdalina* (Bitter Leaf) and *Gongronema latifolium* (Utazi) on the Pancreatic [beta]-Cells of Streptozotocin-Induced Diabetic Rats. *British Journal of Medicine and Medical Research*. 2011, 1(1):24.
- Ademola IO, Eloff JN. Anthelmintic activity of acetone extract and fractions of *Vernonia amygdalina* against *Haemonchus contortus* eggs and larvae. *Tropical Animal Health and Production*. 2011;43(2):521-7.
- Adediran OA, Uwalaka EC. Effectiveness evaluation of levamisole, albendazole, ivermectin, and *Vernonia amygdalina* in West African dwarf goats. *Journal of Parasitology Research*. 2015; 2015.
- Abay SM, Lucantoni L, Dahiya N, Dori G, Dembo EG, Esposito F, Lupidi G, Ogboi S, Ouédraogo RK, Sinisi A, Tagliatella-Scafati O. Plasmodium transmission blocking activities of *Vernonia amygdalina* extracts and isolated compounds. *Malaria Journal*. 2015; 14(1):288.
- Zofou D, Tene M, Ngenemaya MN, Tane P, Titanji VP. *In vitro* antiparasitodal activity and cytotoxicity of extracts of selected medicinal plants used by traditional healers of Western Cameroon. *Malaria Research and Treatment*. 2011;12:2011.
- Omolola AA, Olorunfemi RM, Adetutu AD, Adeniyi SA, Ebenezer OF. Modulatory effect of methanolic extract of *Vernonia amygdalina* (MEVA) on tert-butyl hydroperoxide-induced erythrocyte haemolysis. *Cell Biochemistry and Function*, 2013;31 (7):545-550
- Adesanoye OA, Farombi EO. In vitro Antioxidant Properties of methanolic leaf extract of *Vernonia amygdalina* Del. *Nigerian Journal of Physiological Sciences*. 2014;29 (2):93-101
- Oyeyemi MO, Soetan KO, Akinpelu OB. Sperm characteristics and haemogram of male albino rats (wistar strain) treated with saponin extract from *Vernonia amygdalina* del. asteraceae 110192. *Journal of Cell and Animal Biology*. 2015; 9(3):26-30.
- Momoh MA, Muhamed U, Agboke AA, Akpabio EI, Osonwa UE. Immunological effect of aqueous extract of *Vernonia amygdalina* and a known immune booster called immunace® and their admixtures on HIV/AIDS clients: a comparative study. *Asian Pacific Journal of Tropical Biomedicine*. 2012;2(3):181-4.
- Ezeonu IM, Asuquo AE, Ukwah BN, Ukoha PO. Immunomodulatory properties of prebiotics extracted from *Vernonia amygdalina*. *African Journal of Traditional, Complementary and Alternative Medicines*. 2016;13 (6):11-7.
- Abdulmalik O, Oladapo OO, Bolaji MO. Effect of aqueous extract of *Vernonia amygdalina* on atherosclerosis in rabbits. *ARYA Atherosclerosis*. 2016; 12(1):35.
- Olujimi OO, Onifade ON, Towolawi AT, Akinhanmi TF, Afolabi AA, Olanite KA. Phyto-metals screening of selected anti-diabetic herbs and infused concoctions. *Asian Pacific Journal of Tropical Biomedicine*. 2017;7 (10):909-14.
- 23Nabukenya I, Rubaire-Akiiki C, Mugizi D, Kateregga J, Olila D. Sub-acute toxicity of aqueous extracts of *Tephrosia vogelii*, *Vernonia amygdalina* and *Senna occidentalis* in rats. *Nat Prod Chem Res* 2014;2: 143. doi: 10.4172/2329-6836.1000143